arXiv:math/0309372v3 [math.QA] 4 Apr 2004

IDENTITIES BETWEEN q-HYPERGEOMETRIC AND HYPERGEOMETRIC INTEGRALS OF DIFFERENT DIMENSIONS

V. TARASOV * AND A. VARCHENKO $^{\diamond}$

**St. Petersburg Branch of Steklov Mathematical Institute*
Fontanka 27, St. Petersburg 191011, Russia

$^{\diamond}$*Department of Mathematics, University of North Carolina at Chapel Hill*
Chapel Hill, NC 275993̃250, USA

September 2003

Abstract. Given complex numbers m_1, l_1 and nonnegative integers m_2, l_2, such that $m_1 + m_2 = l_1 + l_2$, for any $a, b = 0, \ldots, \min(m_2, l_2)$ we define an l_2-dimensional Barnes type q-hypergeometric integral $I_{a,b}(z, \mu; m_1, m_2, l_1, l_2)$ and an l_2-dimensional hypergeometric integral $J_{a,b}(z, \mu; m_1, m_2, l_1, l_2)$. The integrals depend on complex parameters z and μ. We show that $I_{a,b}(z, \mu; m_1, m_2, l_1, l_2)$ equals $J_{a,b}(e^{\mu}, z; l_1, l_2, m_1, m_2)$ up to an explicit factor, thus establishing an equality of l_2-dimensional q-hypergeometric and m_2 dimensional hypergeometric integrals. The identity is based on the $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality for the qKZ and dynamical difference equations.

1. Introduction

1.1. q-Hypergeometric integrals.

Let κ be a positive number. Let m_1, l_1 be complex numbers and m_2, l_2 nonnegative integers such that

$$m_1 + m_2 \,=\, l_1 + l_2 \,.$$

We say that an integer a is *admissible* with respect to m_2, l_2 if

$$0 \leqslant a \leqslant \min(m_2, l_2)\,.$$

For a pair of admissible numbers a, b we define a function $I_{a,b}(z, \mu; m_1, m_2, l_1, l_2)$ of complex variables z, μ. The function is defined as an l_2-dimensional Barnes type q-hypergeometric integral:

$$I_{a,b}(z, \mu; m_1, m_2, l_1, l_2) \,= \tag{1.1}$$

$$= \int\limits_{\delta_{l_2}(z; m_1, m_2)} \Phi_{l_2}(t, z, \mu; m_1, m_2)\, w_{l_2-a,a}(t, z; m_1, m_2)\, W_{l_2-b,b}(t, z; m_1, m_2)\, dt^{l_2}\,.$$

Here $t = (t_1, \ldots, t_{l_2})$ and $dt^{l_2} = dt_1 \ldots dt_{l_2}$. The functions $\Phi_{l_2}(t, z, \mu; m_1, m_2)$, $w_{l_2-a,a}(t, z; m_1, m_2)$ and $W_{l_2-b,b}(t, z; m_1, m_2)$ are defined below. The l_2-dimensional integration contour $\delta_{l_2}(z; m_1, m_2)$ lies in $\mathbb{C}^{l_2}$ and is also defined below.

---

*Supported in part by RFFI grant 02˜01˜00085a and CRDF grant RM1˜2334M̃O˜02

E-mail: *vt@pdmi.ras.ru*

$^{\diamond}$Supported in part by NSF grant DMS˜0244579

E-mail: *anv@email.unc.edu*

The *q-master* function Φ_l is defined by the formula

$$\Phi_l(t_1,\ldots,t_l,z,\mu;m_1,m_2) =$$

$$= \exp\bigl(\mu\sum_{u=1}^{l} t_u/\kappa\bigr)\,\prod_{u=1}^{l}\frac{\Gamma(t_u/\kappa)\,\Gamma\bigl((t_u-z)/\kappa\bigr)}{\Gamma\bigl((t_u+m_1)/\kappa\bigr)\,\Gamma\bigl((t_u-z+m_2)/\kappa\bigr)}\prod_{1\leqslant u<v\leqslant l}\frac{\Gamma\bigl((t_u-t_v+1)/\kappa\bigr)}{\Gamma\bigl((t_u-t_v-1)/\kappa\bigr)}\,.$$

The *rational weight* function $w_{l-a,a}$ is defined by the formula

$$w_{l-a,a}(t_1,\ldots,t_l,z;m_1,m_2) = \prod_{1\leqslant u<v\leqslant l}\frac{t_u-t_v}{t_u-t_v-1}\,\prod_{u=1}^{l}\frac{1}{t_u+m_1}\;\times$$

$$\times\;\mathrm{Sym}\Bigl[\,\prod_{u=l-a+1}^{l}\frac{t_u}{t_u-z+m_2}\prod_{1\leqslant u<v\leqslant l}\frac{t_u-t_v-1}{t_u-t_v}\,\Bigr]$$

where $\mathrm{Sym}\,f(t_1,\ldots,t_l) = \sum_{\sigma\in S_l} f(t_{\sigma_1},\ldots,t_{\sigma_l})$. The *trigonometric weight* function $W_{l-b,b}$ is defined by the formula

$$W_{l-b,b}(t_1,\ldots,t_l,z;m_1,m_2) = \prod_{1\leqslant u<v\leqslant l}\frac{\sin\bigl(\pi(t_u-t_v)/\kappa\bigr)}{\sin\bigl(\pi(t_u-t_v-1)/\kappa\bigr)}\,\prod_{u=1}^{l}\frac{e^{-\pi i t_u/\kappa}}{\sin\bigl(\pi(t_u+m_1)/\kappa\bigr)}\;\times$$

$$\times\;\mathrm{Sym}\Bigl[\,\prod_{u=l-b+1}^{l}\frac{e^{\pi i z/\kappa}\sin(\pi t_u/\kappa)}{\sin\bigl(\pi(t_u-z+m_2)/\kappa\bigr)}\prod_{1\leqslant u<v\leqslant l}\frac{\sin\bigl(\pi(t_u-t_v-1)/\kappa\bigr)}{\sin\bigl(\pi(t_u-t_v)/\kappa\bigr)}\,\Bigr].$$

Integral (1.1) is defined for $0<\mathrm{Im}\,\mu<2\pi$. With respect to other parameters we define integral (1.1) by analytic continuation from the region where m_1,m_2 are complex numbers with negative real parts and $\mathrm{Re}\,z=0$. In that case we put

$$\delta_l(z;m_1,m_2) = \{\,(t_1,\ldots,t_l)\in\mathbb{C}^l \mid \mathrm{Re}\,t_u=\varepsilon\,,\;\; u=1,\ldots,l\,\}$$

where ε is a positive number less than $\min(-\mathrm{Re}\,m_1\,,-\mathrm{Re}\,m_2)$. In the considered region of parameters the integrand in (1.1) is well defined on $\delta_{l_2}(z;m_1,m_2)$ for any a,b, the integral is convergent and gives a meromorphic function of z,m_1,m_2, see [TV1]. It is also known that $I_{a,b}(z,\mu;m_1,m_2,l_1,l_2)$ can be analytically continued to a nonnegative integer value of m_2, if a,b are admissible with respect to m_2,l_2 at that point, and the analytic continuation is given by the integral over a suitable deformation of the imaginary plane $\{\,(t_1,\ldots,t_{l_2})\in\mathbb{C}^{l_2} \mid \mathrm{Re}\,t_u=0\,,\; u=1,\ldots,l_2\,\}$, see [MuV].

Remark. There is an alternative way to describe the integrand of integral (1.1), again writing it down as a product of three factors. Namely, consider the functions

$$\Xi_l(t_1,\ldots,t_l,z,\mu;m_1,m_2) = (-\pi\kappa)^{-l(l+3)/2}\exp\bigl((\mu-\pi i)\sum_{u=1}^{l}t_u/\kappa\bigr)\;\times$$

$$\times\prod_{u=1}^{l}\Gamma(t_u/\kappa)\,\Gamma\bigl(-(m_1+t_u)/\kappa\bigr)\,\Gamma\bigl((t_u-z)/\kappa\bigr)\,\Gamma\bigl((z-m_2-t_u)/\kappa\bigr)\;\times$$

$$\times\prod_{1\leqslant u<v\leqslant l}(t_u-t_v)\,\sin\bigl(\pi(t_u-t_v)/\kappa\bigr)\,\Gamma\bigl((t_u-t_v+1)/\kappa\bigr)\,\Gamma\bigl((t_v-t_u+1)/\kappa\bigr)\,,$$

$$p_{l-a,a}(t_1,\dots,t_l,z;m_2)\,=\,\operatorname{Sym}\Biggl[\,\prod_{u=1}^{l-a}(t_u-z+m_2)\prod_{u=l-a+1}^{l}t_u\prod_{1\leqslant u<v\leqslant l}\frac{t_u-t_v-1}{t_u-t_v}\,\Biggr]\,,$$

$$P_{l-b,b}(t_1,\dots,t_l,z;m_2)\,=\,\exp\bigl(\pi ibz/\kappa\bigr)\,\times$$
$$\times\,\operatorname{Sym}\Biggl[\,\prod_{u=1}^{l-b}\sin\bigl(\pi(t_u-z+m_2)/\kappa\bigr)\prod_{u=l-b+1}^{l}\sin(\pi t_u/\kappa)\prod_{1\leqslant u<v\leqslant l}\frac{\sin\bigl(\pi(t_u-t_v-1)/\kappa\bigr)}{\sin\bigl(\pi(t_u-t_v)/\kappa\bigr)}\,\Biggr]\,.$$

Then

$$\Phi_{l_2}(t,z,\mu;m_1,m_2)\,w_{l_2-a,a}(t,z;m_1,m_2)\,W_{l_2-b,b}(t,z;m_1,m_2)\,=$$
$$=\,\Xi_{l_2}(t,z,\mu;m_1,m_2)\,p_{l_2-a,a}(t,z;m_2)\,P_{l_2-b,b}(t,z;m_2)\,.$$

In this description of the integrand the function Ξ_l is such that it contains all the poles of the integrand and has no zeros anywhere but on the shifted diagonals: $(t_u-t_v)\in\kappa\mathbb{Z}$, the functions $p_{l-a,a}$ are polynomials in $t_1,\dots,t_l,z,m_2$, and the functions $P_{l-b,b}$ are trigonometric polynomials in $t_1,\dots,t_l,z,m_2$.

1.2. **Hypergeometric integrals.** Let κ be a positive number. Let m_1,l_1 be complex numbers and m_2,l_2 nonnegative integers such that

$$m_1+m_2\,=\,l_1+l_2\,.$$

For a pair of admissible numbers a,b we define a function $J_{a,b}(z,\mu;m_1,m_2,l_1,l_2)$ of complex variables z,μ. The function is defined as an l_2-dimensional hypergeometric integral:

$$J_{a,b}(z,\mu;m_1,m_2,l_1,l_2)\,=\int\limits_{\gamma_{l_2-b,b}(z)}\Psi_{l_2}(t,z,\mu;m_1,m_2)\,g_{l_2-a,a}(t,z)\,dt^{l_2}\,. \tag{1.2}$$

Here $t=(t_1,\dots,t_{l_2})$, $dt^{l_2}=dt_1\dots dt_{l_2}$. The functions $\Psi_{l_2}(t,z,\mu;m_1,m_2)$ and $g_{l_2-a,a}(t,z)$ are defined below. The l_2-dimensional integration contour $\gamma_{l_2-b,b}(z)$ lies in $\mathbb{C}^{l_2}$ and is also defined below.

The *master* function Ψ_l is defined by the formula

$$\Psi_l(t_1,\dots,t_l,z,\mu;m_1,m_2)\,=$$
$$=\,\prod_{u=1}^{l}t_u^{(\mu+m_1+m_2-2l+1)/\kappa}(1-t_u)^{-m_1/\kappa}(z-t_u)^{-m_2/\kappa}\prod_{1\leqslant u<v\leqslant l}(t_u-t_v)^{2/\kappa}\,.$$

The *weight* function $g_{l-a,a}$ is defined by the formula

$$g_{l-a,a}(t_1,\dots,t_l,z)\,=\,\operatorname{Sym}\Biggl[\,\prod_{u=1}^{l-a}\frac{1}{1-t_u}\prod_{u=l-a+1}^{l}\frac{1}{z-t_u}\,\Biggr]\,.$$

We define the integral in (1.2) by analytic continuation from the region

$$z\neq 0\,,\qquad 0<\arg z<2\pi\,,\qquad \operatorname{Re}\mu\ll 0\,. \tag{1.3}$$

In that region the integration contour $\gamma_{l-b,b}(z)$ is shown in the picture.



Picture 1. Integration contour $\gamma_{l-b,b}(z)$.

It has the form $\gamma_{l-b,b}(z) = \{\,(t_1,\dots,t_l)\in\mathbb{C}^l \mid t_u\in\mathcal{C}_u\,,\ \ u=1,\dots,l\,\}$. Here $\mathcal{C}_u$, $u=1,\dots,l$, are non-intersecting oriented loops in $\mathbb{C}$. The first b loops start at infinity in the direction of z, go around z, and return to infinity in the same direction. For $1\leqslant u<v\leqslant b$ the loop $\mathcal{C}_u$ lies inside the loop $\mathcal{C}_v$. The last $l-b$ loops start at infinity in the real positive direction, go around 1, and return to infinity in the same direction. For $b+1\leqslant u<v\leqslant l$ the loop $\mathcal{C}_u$ lies inside the loop $\mathcal{C}_v$.

If z,μ are in region (1.3), we fix a univalued branch of the master function Ψ_l over $\gamma_{l-b,b}(z)$ by fixing the arguments of all its factors. Namely, we assume that at the point of $\gamma_{l-b,b}(z)$ where all numbers $t_1/z,\dots,t_b/z,t_{b+1},\dots,t_l$ belong to $(0,1)$ we have

$$\arg t_u\in[0,2\pi)\,,\ \ \arg(1-t_u)\in(-\pi,\pi)\,,\ \ \arg(z-t_u)\in(0,2\pi)\,,\ \ \arg(t_u-t_v)\in[0,2\pi)\,,$$

for $u=1,\dots,l$, $v=u+1,\dots,l$. Integral (1.2) is convergent in region (1.3).

1.3. Main result.

Theorem 1. *Let κ be a generic positive number. Let m_1,l_1 be complex numbers and m_2,l_2 nonnegative integers, such that $m_1+m_2=l_1+l_2$. Let $0<\operatorname{Im}\mu<2\pi$. Then for any $a,b=1,\dots,\min(m_2,l_2)$ we have*

$$C_b(m_1,m_2,l_1,l_2)\,I_{a,b}(z,\mu;m_1,m_2,l_1,l_2)\;= \tag{1.4}$$

$$=\;D_b(m_1,m_2,l_1,l_2)\,E_b(l_1,l_2)\,X(z;m_1,m_2)\,Y(\mu;m_1,m_2,l_1,l_2)\,J_{a,b}(e^\mu,z;l_1,l_2,m_1,m_2)$$

where

$$C_b(m_1,m_2,l_1,l_2)\;=\;\frac{(2\pi i)^{-l_2}}{l_2!\,(l_2-b)!\,b!}\prod_{j=0}^{l_2-b-1}\sin\bigl(\pi(m_1-j)/\kappa\bigr)\prod_{j=0}^{b-1}\sin\bigl(\pi(m_2-j)/\kappa\bigr)\;\times$$

$$\times\prod_{j=0}^{l_2-1}\frac{\Gamma(1+1/k)\,\Gamma\bigl(1+(m_1-j)/\kappa\bigr)}{\Gamma\bigl(1+(j+1)/\kappa\bigr)}\;,$$

$$D_b(m_1,m_2,l_1,l_2)\;=\;(2i)^{-m_2}\prod_{j=0}^{m_2-b-1}\frac{1}{\sin\bigl(\pi(j+1)/\kappa\bigr)}\prod_{j=0}^{b-1}\frac{1}{\sin\bigl(\pi(j+1)/\kappa\bigr)}\;\times$$

$$\times\prod_{j=0}^{m_2-1}\frac{\Gamma\bigl(1+(l_1-j)/\kappa\bigr)}{\Gamma(-1/\kappa)\,\Gamma\bigl(1+(j+1)/\kappa\bigr)}\;,$$

$$E_b(l_1, l_2) = \exp\bigl(\pi i (b^2 - (b - l_2)(l_1 + l_2) - l_2(l_2 - 1)/2)/\kappa\bigr)\,,$$

$$X(t; m_1, m_2) = \prod_{j=0}^{m_2-1} \frac{\Gamma\bigl((j - m_1 - t)/\kappa\bigr)}{\Gamma\bigl((j + 1 - t))/\kappa\bigr)}\,, \tag{1.5}$$

$$Y(\mu; m_1, m_2, l_1, l_2) = e^{\mu l_2(l_2 - 2m_1 - 1)/2\kappa}\,(1 - e^{\mu})^{(l_1+1)l_2/\kappa}\,, \quad \arg(1 - e^{\mu}) \in (-\pi, \pi)\,. \tag{1.6}$$

Remark. There is a similar theorem establishing an equality of suitable hypergeometric integrals of different dimensions, see [TV4]. The factor $D_b(m_1, m_2, l_1, l_2)$ in the present paper corresponds to the normalization factor $C_b(l_1, l_2, m_1, m_2)$ in [TV4] and contains the same product of sines and gamma-functions.

Example. Let $m_2 = l_2 = 1$. In this case formula (1.4) becomes the classical equality of two integral representations of the Gauss hypergeometric function ${}_2F_1$. For instance, if $a = b = 0$, then taking $\alpha = -m_1/\kappa$, $\beta = -(z + m_1)/\kappa$, $\gamma = (1 - z - m_1)/\kappa$, after simple transformations one gets:

$$\frac{1}{2\pi i} \int\limits_{-i\infty-\varepsilon}^{+i\infty-\varepsilon} e^{s(\mu - \pi i)}\,\Gamma(-s)\,\Gamma(s + \alpha)\,\frac{\Gamma(s + \beta)}{\Gamma(s + \gamma)}\,ds =$$

$$= (1 - e^{\mu})^{\gamma-\alpha-\beta}\,\frac{\Gamma(\beta)}{\Gamma(\gamma - \alpha)} \int\limits_1^{+\infty} t^{\beta}\,(t - 1)^{\alpha-1}\,(t - e^{\mu})^{\beta-\gamma}\,dt =$$

$$= \frac{\Gamma(\beta)}{\Gamma(\gamma - \alpha)} \int\limits_0^1 u^{\alpha-1}\,(1 - u)^{\gamma-\alpha-1}\,(1 - ue^{\mu})^{-\beta}\,du = \frac{\Gamma(\alpha)\,\Gamma(\beta)}{\Gamma(\gamma)}\,{}_2F_1(\alpha, \beta; \gamma; e^{\mu})\,.$$

Here it is assumed that $\operatorname{Re}\gamma > \operatorname{Re}\alpha > 0$, $\operatorname{Re}\beta > 0$ and $0 < \varepsilon < \min(\operatorname{Re}\alpha, \operatorname{Re}\beta)$. The second equality is obtained by the change of integration variable $u = (t - 1)/(t - e^{\mu})$.

Theorem 1 claims that an l_2-dimensional q-hypergeometric integral equals an m_2 dimensional hypergeometric integral up to an explicit factor. Note that in the first integral the numbers m_1, m_2 are shifts of arguments of the gamma-functions entering its q-master function, while in the second integral m_2 is its dimension and m_1 is not present explicitly.

It is well known that studying asymptotics of integrals with respect to their dimension is an interesting problem appearing, for instance, in the theory of orthogonal polynomials and in matrix models. The duality of the theorem allows us to study asymptotics of integrals with respect to their dimension. Namely, assume that in the 4-tuple (m_1, m_2, l_1, l_2) the nonnegative integer l_2 tends to infinity while the numbers m_1 and m_2 remain fixed. Then $I_{a,b}(z, \mu; m_1, m_2, l_1, l_2)$ is a q-hypergeometric integral of growing dimension l_2, whose master function has fixed shifts m_1, m_2. At the same time, $J_{a,b}(e^{\mu}, z; l_1, l_2, m_1, m_2)$ is a hypergeometric integral of the fixed dimension m_2, whose master function has growing exponents l_1, l_2. The asymptotics of $J_{a,b}(e^{\mu}, z; l_1, l_2, m_1, m_2)$ can be calculated using the steepest descent method. An example of such calculation is given in [TV4].

Similarly one can assume that in the 4-tuple (m_1, m_2, l_1, l_2) the nonnegative integer m_2 tends to infinity while the numbers l_1 and l_2 are fixed. Then $J_{a,b}(e^{\mu}, z; l_1, l_2, m_1, m_2)$ is a hypergeometric integral of growing dimension m_2, whose master function has fixed exponents l_1, l_2. On the other hand, $I_{a,b}(z, \mu; m_1, m_2, l_1, l_2)$ is a q-hypergeometric integral of the fixed dimension l_2, whose master function has growing shifts m_1, m_2. The asymptotics of $I_{a,b}(z, \mu; m_1, m_2, l_1, l_2)$ in principle can be calculated using the Stirling formula for asymptotics of the gamma-function.

To prove Theorem 1 we show that the matrices:

$$\bigl(I_{a,b}(z, \mu; m_1, m_2, l_1, l_2)\bigr)_{0 \leqslant a,b \leqslant \min(m_2, l_2)} \tag{1.7}$$

and

$$X(z; m_1, m_2)\,\bigl(J_{a,b}(e^{\mu}, z; l_1, l_2, m_1, m_2)\bigr)_{0 \leqslant a,b \leqslant \min(m_2, l_2)}$$

satisfy the same system of first order linear difference equations with respect to z, see Corollary 4 and Theorems 5, 7. Studying asymptotics of those matrices as $\operatorname{Re} z \to -\infty$ allows us to compute the connection matrix, thus proving Theorem 1.

The fact that both matrices satisfy the same system of difference equations is based on the duality of the *qKZ* and dynamical equations for $\mathfrak{gl}_k$ and $\mathfrak{gl}_n$ [TV3], which is a generalization of the duality between the rational differential *KZ* and dynamical equations observed in [TL]. Namely, in [TV2] a system of dynamical difference equations was introduced. It is proved there that the dynamical difference equations are compatible with the trigonometric *KZ* differential equations. In [MV] hypergeometric solutions of the trigonometric *KZ* and dynamical difference equations were presented. On the other hand, q-hypergeometric solutions of the rational *qKZ* difference equations equations were constructed in [TV1]. It was shown in [TV3] that the system of dynamical difference equations for $\mathfrak{gl}_k$ and the system of rational *qKZ* equations for $\mathfrak{gl}_n$ are naturally transformed into each other under the $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality. In this way one gets two sets of solutions of the same system of equations, and in principle, these two sets of solutions can be identified. Theorem 1 is a realization of this idea for the case of $k = n = 2$. We will discuss the case of an arbitrary pair k, n in a separate paper.

Let us make an additional remark. In [TV3] it was introduced a system of dynamical differential equations which is transformed under the $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality to the trigonometric *KZ* differential equations. It is proved in [TV5] that the dynamical differential equations are compatible with the rational *qKZ* difference equations. It is shown in [TV6] that the q-hypergeometric solutions of the *qKZ* equations satisfy the dynamical differential equations. In the present case of $k = n = 2$ this means that the matrices (1.7) and

$$Y(\mu; m_1, m_2, l_1, l_2)\,\bigl(J_{a,b}(e^{\mu}, z; l_1, l_2, m_1, m_2)\bigr)_{0 \leqslant a,b \leqslant \min(m_2, l_2)} \tag{1.8}$$

satisfy the same system of first order linear differential equations with respect to μ. Therefore, the connection matrix of (1.7) and (1.8) does not depend on μ, which agrees with formula (1.4). Other factors in (1.4) also have natural explanation in terms of the q-deformation of the $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality.

The rest of the paper is as follows. In Section 2 we discuss the $(\mathfrak{gl}_2, \mathfrak{gl}_2)$ duality for the *qKZ* and dynamical equations. In Section 3 we describe their hypergeometric solutions and prove Theorem 1. In Section 4 we give some facts about the dynamical differential equations.

The authors thank Y. Markov for useful discussions.

2. The $(\mathfrak{gl}_2, \mathfrak{gl}_2)$ duality for KZ and dynamical equations

In this section we follow the exposition in [TV3].

2.1. **The trigonometric KZ and associated dynamical difference equations.** Let E_{ij}, $i, j = 1, 2$, be the standard generators of the Lie algebra $\mathfrak{gl}_2$. Let $\mathfrak{h} = \mathbb{C}\cdot E_{11} \oplus \mathbb{C}\cdot E_{22}$ be the Cartan subalgebra, while E_{12} and E_{21} are respectively the positive and negative root vectors.

The trigonometric r-matrix $r(z) \in \mathfrak{gl}_2^{\otimes 2}$ is defined by the formula

$$r(z) = \frac{1}{z-1}\bigl((z+1)(E_{11}\otimes E_{11} + E_{22}\otimes E_{22})/2 + zE_{12}\otimes E_{21} + E_{21}\otimes E_{12}\bigr)$$

Fix a non-zero complex number κ. The *trigonometric Knizhnik-Zamolodchikov* (KZ) *operators* $\nabla_1, \dots, \nabla_n$ are the following differential operators with coefficients in $U(\mathfrak{gl}_2)^{\otimes n}$ acting on functions of complex variables $z_1, \dots, z_n, \lambda_1, \lambda_2$:

$$\nabla_a(z_1, \dots, z_n, \lambda_1, \lambda_2) = \kappa z_a \frac{\partial}{\partial z_a} - \sum_{i=1}^{2}\bigl(\lambda_i - \frac{1}{2}\sum_{b=1}^{n} E_{ii}^{(b)}\bigr) E_{ii}^{(a)} - \sum_{\substack{b=1 \\ b\neq a}}^{n} r(z_a/z_b)^{(ab)} . \tag{2.1}$$

Here $E_{ii}^{(b)} = 1 \otimes \dots \otimes \underset{b\text{-th}}{E_{ii}} \otimes \dots \otimes 1$, and the meaning of $r(z_a/z_b)^{(ab)}$ is similar. It is known that the operators $\nabla_1, \dots, \nabla_n$ pairwise commute.

Let $V_1, \dots, V_n$ be $\mathfrak{gl}_2$-modules. The *trigonometric KZ equations* for a function $U(z_1, \dots, z_n, \lambda_1, \lambda_2)$ with values in $V_1 \otimes \dots \otimes V_n$ are

$$\nabla_a(z_1, \dots, z_n, \lambda_1, \lambda_2)\, U(z_1, \dots, z_n, \lambda_1, \lambda_2) = 0 , \qquad a = 1, \dots, n . \tag{2.2}$$

Let T_u be the difference operator acting on functions $f(u)$ by the formula

$$(T_u f)(u) = f(u + \kappa) .$$

Introduce a series $B(t)$ depending on a complex variable t:

$$B(t) = 1 + \sum_{s=1}^{\infty} E_{21}^s E_{12}^s \prod_{j=1}^{s} \frac{1}{j\,(t - E_{11} + E_{22} - j)} .$$

For any $\mathfrak{gl}_2$-module V with a locally nilpotent action of E_{12} and finite-dimensional weight subspaces the series $B(t)$ has a well-defined action in any weight subspace $V[\mu] \subset V$ as a rational $\operatorname{End}\bigl(V[\mu]\bigr)$-valued function of t.

Let $V_1, \dots, V_n$ be $\mathfrak{gl}_2$-modules as above. Introduce the *dynamical difference operators* Q_1, Q_2 acting on $V_1 \otimes \dots \otimes V_n$-valued functions of complex variables $z_1, \dots, z_n, \lambda_1, \lambda_2$ by the formulae

$$Q_1(z_1, \dots, z_n, \lambda_1, \lambda_2) = \bigl(B(\lambda_1 - \lambda_2)\bigr)^{-1} \prod_{a=1}^{n} z_a^{-E_{11}^{(a)}}\, T_{\lambda_1} , \tag{2.3}$$

$$Q_2(z_1, \dots, z_n, \lambda_1, \lambda_2) = \prod_{a=1}^{n} z_a^{-E_{22}^{(a)}}\, B(\lambda_1 - \lambda_2 - \kappa)\, T_{\lambda_2} .$$

One can see that the operators Q_1, Q_2 commute.

Proposition 2 ([TV2]). *One has* $[\nabla_a, Q_i] = 0$ *for all* $a = 1, \dots, n$ *and* $i = 1, 2$.

The *dynamical difference equations* associated with the trigonometric KZ equations for a function $U(z_1,\dots,z_n,\lambda_1,\lambda_2)$ with values in $V_1\otimes\dots\otimes V_n$ are

$$Q_i(z_1,\dots,z_n,\lambda_1,\lambda_2)\,U(z_1,\dots,z_n,\lambda_1,\lambda_2)\,=\,U(z_1,\dots,z_n,\lambda_1,\lambda_2)\,,\qquad i=1,2\,. \tag{2.4}$$

The trigonometric KZ and dynamical difference operators preserve the weight decomposition of $V_1\otimes\dots\otimes V_n$. Thus the KZ and dynamical equations can be considered as equations for a function $U(z_1,\dots,z_n,\lambda_1,\lambda_2)$ taking values in a given weight subspace of $V_1\otimes\dots\otimes V_n$.

2.2. **The** *qKZ* **equations.** Let V,W be irreducible highest weight $\mathfrak{gl}_2$-modules with highest weight vectors v,w, respectively. There is a unique rational function $R_{VW}(t)$ taking values in $\mathrm{End}\,(V\otimes W)$ such that

$$\bigl[R_{VW}(t)\,,\,g\otimes 1+1\otimes g\bigr]\,=\,0\qquad\text{for any}\quad g\in\mathfrak{gl}_2\,,$$

$$R_{VW}(t)\,\bigl(E_{21}\otimes E_{11}+E_{22}\otimes E_{21}+t\,E_{21}\otimes 1\bigr)\,=\,\bigl(E_{11}\otimes E_{21}+E_{21}\otimes E_{22}+t\,E_{21}\otimes 1\bigr)\,R_{VW}(t)\,,$$

$$R_{VW}(t)\,v\otimes w\,=\,v\otimes w\,.$$

The function $R_{VW}(t)$ is called the rational R-matrix for the tensor product $V\otimes W$. It comes from the representation theory of the Yangian $Y(\mathfrak{gl}_2)$.

Let $V_1,\dots,V_n$ be irreducible highest weight $\mathfrak{gl}_2$-modules. Introduce the *qKZ difference operators* $Z_1,\dots,Z_n$ acting on $V_1\otimes\dots\otimes V_n$-valued functions of complex variables $z_1,\dots,z_n,\lambda_1,\lambda_2$ by the formula

$$\begin{aligned} Z_a(z_1,\dots,z_n,\lambda_1,\lambda_2)\,&=\,\bigl(R_{an}(z_a-z_n)\dots R_{a,a+1}(z_a-z_{a+1})\bigr)^{-1}\times\\ &\times\,\lambda_1^{-E_{11}^{(a)}}\lambda_2^{-E_{22}^{(a)}}\,R_{1a}(z_1-z_a-\kappa)\dots R_{a-1,a}(z_{a-1}-z_a-\kappa)\,T_{z_a}\,. \end{aligned} \tag{2.5}$$

These operators are called the *qKZ operators*. It is known that they pairwise commute [FR]. The difference equations

$$Z_a(z_1,\dots,z_n,\lambda_1,\lambda_2)\,U(z_1,\dots,z_n,\lambda_1,\lambda_2)\,=\,U(z_1,\dots,z_n,\lambda_1,\lambda_2)\,,\qquad a=1,\dots,n\,, \tag{2.6}$$

for a $V_1\otimes\dots\otimes V_n$-valued function $U(z_1,\dots,z_n,\lambda_1,\lambda_2)$ are called the *qKZ equations*.

The *qKZ* operators preserve the weight decomposition of $V_1\otimes\dots\otimes V_n$. Thus the *qKZ* and dynamical differential equations can be considered as equations for a function $U(z_1,\dots,z_n,\lambda_1,\lambda_2)$ taking values in a given weight subspace of $V_1\otimes\dots\otimes V_n$.

2.3. **The duality.** For a complex number m, denote M_m the Verma module over $\mathfrak{gl}_2$ with highest weight $(m,0)$ and highest weight vector v_m. The vectors $E_{2,1}^d v_m$, $d\in\mathbb{Z}_{\geqslant 0}$, form a basis in M_m.

For a non-negative integer m, denote L_m the irreducible $\mathfrak{gl}_2$-module with highest weight $(m,0)$ and highest weight vector v_m. The vectors $E_{2,1}^d v_m$, $d=0,\dots,m$, form a basis in L_m.

Let m_1, l_1 be complex numbers and m_2, l_2 non-negative integers such that $m_1 + m_2 = l_1 + l_2$. Consider the weight subspace $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ of the tensor product $M_{m_1} \otimes L_{m_2}$. The weight subspace has a basis

$$F^a(m_1, m_2, l_1, l_2) \,=\, \frac{1}{(l_2 - a)!\, a!}\, E_{2,1}^{l_2 - a} v_{m_1} \otimes E_{2,1}^{a} v_{m_2}\,, \qquad a = 0, \dots, \min(m_2, l_2)\,. \tag{2.7}$$

There is a linear isomorphism

$$\begin{aligned} \varphi\,:\,(M_{m_1} \otimes L_{m_2})[l_1, l_2] \,&\to\, (M_{l_1} \otimes L_{l_2})[m_1, m_2]\,, \\ F^a(m_1, m_2, l_1, l_2) \,&\mapsto\, F^a(l_1, l_2, m_1, m_2)\,. \end{aligned} \tag{2.8}$$

Theorem 3 ([TV3]). *The isomorphism φ transforms the qKZ operators acting in $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ into the dynamical difference operators acting in $(M_{l_1} \otimes L_{l_2})[m_1, m_2]$. More precisely, we have*

$$\varphi\, Z_1(z_1, z_2, \lambda_1, \lambda_2) \,=\, \bigl(G(z_1 - z_2; m_1, m_2)\bigr)^{-1} Q_1(\lambda_1, \lambda_2, z_1, z_2)\, \varphi\,,$$

$$\varphi\, Z_2(z_1, z_2, \lambda_1, \lambda_2) \,=\, G(z_1 - z_2 - \kappa; m_1, m_2)\, Q_2(\lambda_1, \lambda_2, z_1, z_2)\, \varphi\,,$$

where

$$G(t; m_1, m_2) \,=\, \prod_{j=0}^{m_2 - 1} \frac{t + j - m_1}{t + j + 1}\,.$$

Let $S(t)$ be any solution of the equation

$$S(t + \kappa) \,=\, G(t; m_1, m_2)\, S(t)\,.$$

For instance, one of solutions is given by $S(t) = X(-t; m_1, m_2)$, cf. (1.5).

Corollary 4. *Let an $(M_{l_1} \otimes L_{l_2})[m_1, m_2]$-valued function $U(z_1, z_2, \lambda_1, \lambda_2)$ solve the dynamical difference equations*:

$$Q_a(z_1, z_2, \lambda_1, \lambda_2)\, U(z_1, z_2, \lambda_1, \lambda_2) \,=\, U(z_1, z_2, \lambda_1, \lambda_2)\,, \qquad a = 1, 2\,.$$

Then the $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$-valued function

$$\widetilde{U}(z_1, z_2, \lambda_1, \lambda_2) \,=\, S(z_1 - z_2)\, \varphi^{-1}\bigl(U(\lambda_1, \lambda_2, z_1, z_2)\bigr)$$

solves the qKZ equations:

$$Z_a(z_1, z_2, \lambda_1, \lambda_2)\, \widetilde{U}(z_1, z_2, \lambda_1, \lambda_2) \,=\, \widetilde{U}(z_1, z_2, \lambda_1, \lambda_2)\,, \qquad a = 1, 2\,.$$

More facts on the $(\mathfrak{gl}_2, \mathfrak{gl}_2)$ duality for *KZ* and dynamical equations are given in Section 4.

3. Hypergeometric solutions

3.1. **Hypergeometric solutions of the** qKZ **equations.** For any $b = 0, \dots, \min(m_2, l_2)$ define an $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$-valued function

$$\bar{I}_b(z, \mu; m_1, m_2, l_1, l_2) \,=\, \sum_{a=0}^{l_2} I_{a,b}(z, \mu; m_1, m_2, l_1, l_2)\, \frac{1}{(l_2 - a)!\, a!}\, E_{2,1}^{l_2 - a} v_{m_1} \otimes E_{2,1}^{a} v_{m_2}\,.$$

Here the actual range of summation is until $a = \min(m_2, l_2)$, since $E_{2,1}^a v_{m_2} = 0$ for $a > m_2$.

Theorem 5. *Let* $\lambda_i = e^{\mu_i}$, $i = 1, 2$. *For any* $b = 0, \dots, \min(m_2, l_2)$ *the function*

$$U_b(z_1, z_2, \lambda_1, \lambda_2) \,=\, e^{(\mu_1(m_1 z_1 + m_2 z_2 - (m_1^2 + m_2^2)/2) - (\mu_1 - \mu_2)(l_2 z_1 + l_2/2))/\kappa} \,\times \\ \times\, (1 - e^{\mu_2 - \mu_1})^{-l_2/\kappa}\, \bar I_b(z_2 - z_1, \mu_2 - \mu_1; m_1, m_2, l_1, l_2) \tag{3.1}$$

is a solution of the qKZ equations (2.6) *with values in* $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$. *Moreover, if* λ_1/λ_2 *is not real, than any solution of that qKZ equations is a linear combination of functions* $U_b(z_1, z_2, \lambda_1, \lambda_2)$ *with coefficients being* κ*-periodic functions of* z_1, z_2.

The theorem is a direct corollary of the construction of q-hypergeometric solutions of the qKZ equations given in [TV1], [MuV].

We describe asymptotics of the integral $I_{a,b}(z, \mu; m_1, m_2, l_1, l_2)$ as $\operatorname{Re} z \to -\infty$ and μ is fixed. For a positive number κ, complex numbers m, μ, $\operatorname{Im} \mu \in (0, 2\pi)$, and a nonnegative integer l consider the Selberg-type integral

$$A_l(\mu; m) \,=\, \int\limits_{\delta_l(m)} \exp\bigl((\mu - \pi i) \sum_{u=1}^{l} s_u\bigr) \prod_{u=1}^{l} \Gamma(s_u)\, \Gamma(-s_u - m/\kappa) \,\times \\ \times \prod_{\substack{u,v=1 \\ u \neq v}}^{l} \frac{\Gamma(s_u - s_v + 1/\kappa)}{\Gamma(s_u - s_v)}\, ds^l\,. \tag{3.2}$$

The integral is defined by analytic continuation from the region where $\operatorname{Re} m$ is negative. In that case

$$\delta_l(m) \,=\, \{\, (s_1, \dots, s_l) \in \mathbb{C}^l \mid \ \operatorname{Re} s_u = \operatorname{Re} m/2\,, \ \ u = 1, \dots, l \,\}\,.$$

In the considered region of parameters the integrand in (3.2) is well defined on $\delta_l(m)$ and the integral is convergent, see [TV1]. The formula for $A_l(m)$ is well known,

$$A_l(\mu; m) \,= \\ =\, (2\pi i)^l\, e^{(\mu - \pi i)(l - 1 - 2m)l/2\kappa}\, (1 - e^{\mu})^{l(m - l + 1)/\kappa} \prod_{j=0}^{l-1} \frac{\Gamma\bigl(1 + (j+1)/\kappa\bigr)}{\Gamma(1 + 1/\kappa)}\, \Gamma\bigl((j - m)/\kappa\bigr)\,,$$

where $\arg(1 - e^{\mu}) \in (-\pi, \pi)$, see, for example, [TV1].

Remark. Other versions of Selberg-type integrals see in [FSV] and [TV7].

Lemma 6. *Let* $\operatorname{Re} z \to -\infty$ *and* μ *is fixed. Then*

$$I_{a,b}(z, \mu; m_1, m_2, l_1, l_2) \,=\, l_2!\, (l_2 - b)!\, b!\, (-\pi)^{-l_2} \exp(\mu z b/\kappa) \,\times \\ \times\, (-z/\kappa)^{-(2b^2 + b(m_1 - m_2 - 2l_2) + m_2 l_2)/\kappa} A_{l_2 - b}(\mu; m_1)\, A_b(\mu; m_2) \bigl(\delta_{ab} + O(z^{-1})\bigr)\,.$$

The lemma follows from [TV1].

3.2. **Hypergeometric solutions of the trigonometric** ***KZ*** **and difference dynamical equations.** For $b=0,\dots,\min(m_2,l_2)$, define an $M_{l_1}\otimes L_{l_2}[m_1,m_2]$-valued function

$$\bar J_b(z,\mu;l_1,l_2,m_1,m_2) \,=\, \sum_{a=0}^{m_2} J_{a,b}(z,\mu;l_1,l_2,m_1,m_2)\,\frac{1}{(m_2-a)!\,a!}\,E_{2,1}^{m_2-a}v_{l_1}\otimes E_{2,1}^{a}v_{l_2}\,.$$

Here the actual range of summation is until $a=\min(m_2,l_2)$, since $E_{2,1}^{a}v_{l_2}=0$ for $a>l_2$.

Theorem 7. *For any* $b=0,\dots,\min(m_2,l_2)$ *the function*

$$U_b(z_1,z_2,\lambda_1,\lambda_2) \,=\, z_1^{(\lambda_1(l_1-m_2)+\lambda_2 m_2-m_2^2-m_1l_1+l_1^2/2)/\kappa}\,z_2^{l_2(\lambda_1-m_1+l_2/2)/\kappa}\,\times$$
$$\times\,(z_1-z_2)^{l_1l_2/\kappa}\,\bar J_b(z_2/z_1,\lambda_2-\lambda_1;l_1,l_2,m_1,m_2)$$

is a solution of the KZ equations (2.2) *and difference dynamical equations* (2.4) *with values in* $(M_{l_1}\otimes L_{l_2})[m_1,m_2]$.

The theorem is a direct corollary of [MV].

We describe asymptotics of the integral $J_{a,b}(e^{\mu},z;l_1,l_2,m_1,m_2)$ as $\operatorname{Re} z\to-\infty$ and μ is fixed. For a positive number κ, a complex number l and a nonnegative integer m consider the Selberg-type integral

$$B_m(l) \,=\, \int\limits_{\gamma_m} e^{-\sum\limits_{u=1}^{m}s_u/\kappa}\,\prod_{u=1}^{m}(-s_u)^{-1-l/\kappa}\prod_{1\leqslant u<v\leqslant m}(s_u-s_v)^{2/\kappa}\,ds^m\,.$$

The integration contour γ_m has the form

$$\gamma_m \,=\, \{\,(s_1,\dots,s_m)\in\mathbb{C}^m \mid s_u\in\mathcal{C}_u\,,\ u=1,\dots,m\,\}\,,$$

see the picture.



Picture 2. The contour γ_m.

Here $\mathcal{C}_u$, $u=1,\dots,m$, are non-intersecting oriented loops in $\mathbb{C}$. The loops start at $+\infty$, go around 0, and return to $+\infty$. For $u<v$ the loop $\mathcal{C}_u$ lies inside the loop $\mathcal{C}_v$. We fix a univalued branch of the integrand by assuming that at the point of γ_m where all numbers $s_1,\dots,s_m$ are negative we have $\arg(-s_u)=0$ for $u=1,\dots,m$, and $\arg(s_u-s_v)=0$ for $1\leqslant u<v\leqslant m$.

The formula for $B_m(l)$ is well known:

$$B_m(l) \,=\, (-2\pi i)^m\,\kappa^{m(m-1-l)/\kappa}\prod_{j=0}^{m-1}\frac{\Gamma(1-1/\kappa)}{\Gamma(1+(l-j)/\kappa)\,\Gamma(1-(j+1)/\kappa)}\,, \tag{3.3}$$

for example, cf. [TV2], [MTV].

Lemma 8. *Let* $\mathrm{Re}\, z \to -\infty$ *and* μ *is fixed. Then*

$$\begin{aligned} J_{a,b}(e^{\mu}, z; l_1, l_2, m_1, m_2) \,=\, (m_2-b)!\, b!\, e^{\pi i (m_2-b)(2b-l_2)/\kappa} \,\times \\ \times\, e^{\mu b(z+l_1-2m_2+b)/\kappa}\, (1-e^{\mu})^{-(2b^2+b(l_1-l_2-2m_2)+m_2 l_2)/\kappa} \,\times \\ \times\, (-z)^{-(2b^2+b(l_1-l_2-2m_2)+m_2(m_2-l_1-1))/\kappa} B_{m_2-b}(l_1)\, B_b(l_2)\bigl(\delta_{ab}+O(z^{-1})\bigr)\,. \end{aligned}$$

The proof is straightforward.

3.3. **Proof of Theorem 1.** Theorems 5 and 7, and Corollary 4 imply that for any $b = 0, \ldots, \min(m_2, l_2)$ the functions $\bar I_b(z, \mu; m_1, m_2, l_1, l_2)$ and $X(z; m_1, m_2)\, \varphi^{-1}\bigl(\bar J_b(e^{\mu}, z; l_1, l_2, m_1, m_2)\bigr)$ satisfy the same first order difference equation with respect to z with step κ. Hence, for any $a = 0, \ldots, \min(m_2, l_2)$ one has

$$X(z; m_1, m_2)\, J_{a,b}(e^{\mu}, z; l_1, l_2, m_1, m_2) \,=\, \sum_{c=0}^{\min(m_2,l_2)} I_{a,c}(z, \mu; m_1, m_2, l_1, l_2)\, G_{b,c}(z, \mu; m_1, m_2, l_1, l_2)\,, \tag{3.4}$$

the connection coefficients $G_{b,c}(z, \mu; m_1, m_2, l_1, l_2)$ being κ-periodic functions of z and holomorphic functions of μ in the strip $0 < \mathrm{Im}\, \mu < 2\pi$. Taking into account asymptotics of the integrals $I_{a,c}(z, \mu; m_1, m_2, l_1, l_2)$ and $J_{a,b}(e^{\mu}, z; l_1, l_2, m_1, m_2)$ as $\mathrm{Re}\, z \to -\infty$ and μ is fixed, see Lemmas 6 and 8, one can compute the connection coefficients and obtain formula (1.4). Theorem 1 is proved. □

Remark. One can see from formula (1.4) that all connection coefficients $G_{b,c}(z, \mu; m_1, m_2, l_1, l_2)$ in (3.4) as functions of μ are proportional to the same function $Y(\mu; m_1, m_2, l_1, l_2)$. This fact, which is pure computational in the given proof of Theorem 1, can be observed independently in advance, because Theorems 7, 12, and Corollary 11 imply that the functions $\bar I_b(z, \mu; m_1, m_2, l_1, l_2)$ and $Y(\mu; m_1, m_2, l_1, l_2)\, \varphi^{-1}\bigl(\bar J_b(e^{\mu}, z; l_1, l_2, m_1, m_2)\bigr)$ satisfy the same first order differential equation with respect to μ.

4. DIFFERENTIAL DYNAMICAL OPERATORS

Introduce the *dynamical differential operators* D_1, D_2 with coefficients in $U(\mathfrak{gl}_2)^{\otimes n}$ acting on functions of complex variables $z_1, \ldots, z_n, \lambda_1, \lambda_2$ by the formula

$$\begin{aligned} D_i(z_1, \ldots, z_n, \lambda_1, \lambda_2) \,=\, \\ =\, \kappa \lambda_i \frac{\partial}{\partial \lambda_i} + \frac{\widetilde E_{ii}^2}{2} - \sum_{a=1}^{n} z_a E_{ii}^{(a)} - \sum_{j=1}^{2} \sum_{1 \leqslant a < b \leqslant n} E_{ij}^{(a)} E_{ji}^{(b)} - \frac{\lambda_{i'}}{\lambda_i - \lambda_{i'}} (\widetilde E_{21} \widetilde E_{12} - \widetilde E_{22})\,. \end{aligned}$$

Here $\widetilde E_{kl} = \sum_{a=1}^{n} E_{kl}^{(a)}$, and i' is supplementary to i, that is, $\{i, i'\} = \{1, 2\}$.

Proposition 9 ([TV5])**.** *One has* $[Z_a, D_i] = 0$ *for all* $a = 1, \ldots, n$ *and* $i = 1, 2$*, where* $Z_1, \ldots, Z_n$ *are the qKZ operators* (2.5)*.*

The differential equations

$$D_i(z_1, \ldots, z_n, \lambda_1, \lambda_2)\, U(z_1, \ldots, z_n, \lambda_1, \lambda_2) \,=\, 0\,, \qquad i = 1, 2\,, \tag{4.1}$$

for a $V_1 \otimes \ldots \otimes V_n$-valued function $U(z_1, \ldots, z_n, \lambda_1, \lambda_2)$ are called the *dynamical differential equations* associated with the qKZ equations.

Theorem 10 ([TV3]). *The isomorphism* φ, *see* (2.8), *transforms the dynamical differential operators acting in* $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ *into the trigonometric KZ operators* (2.1) *acting in* $(M_{l_1} \otimes L_{l_2})[m_1, m_2]$:

$$\varphi\, D_a(z_1, z_2, \lambda_1, \lambda_2) \,=\, \nabla_a(\lambda_1, \lambda_2, z_1, z_2)\, \varphi\,, \qquad a = 1, 2\,.$$

Corollary 11. *Let an* $(M_{l_1} \otimes L_{l_2})[m_1, m_2]$*-valued function* $U(z_1, z_2, \lambda_1, \lambda_2)$ *solve the trigonometric KZ equations*:

$$\nabla_a(z_1, z_2, \lambda_1, \lambda_2)\, U(z_1, z_2, \lambda_1, \lambda_2) \,=\, 0\,, \qquad a = 1, 2\,.$$

Then the $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$*-valued function*

$$\widetilde{U}(z_1, z_2, \lambda_1, \lambda_2) \,=\, \varphi^{-1}\bigl(U(\lambda_1, \lambda_2, z_1, z_2)\bigr)$$

solves the system of the dynamical differential equations (4.1):

$$D_a(z_1, z_2, \lambda_1, \lambda_2)\, \widetilde{U}(z_1, z_2, \lambda_1, \lambda_2) \,=\, 0\,, \qquad a = 1, 2\,.$$

The next statement describes q-hypergeometric solutions of the dynamical differential equations.

Theorem 12. *For any* $b = 0, \ldots, \min(m_2, l_2)$ *the function* $U_b(z_1, z_2, \lambda_1, \lambda_2)$ *defined by* (3.1) *is a solution of equations* (4.1) *with values in* $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$.

The theorem follows from [TV6].

References

[FR] Frenkel. I. B., Reshetikhin N. Yu.,: Quantum affine algebras and holonomic difference equations, Comm. Math. Phys. **146** (1992), 1–60.

[FSV] Felder, G., Stevens, L., Varchenko, A.: Elliptic Selberg integrals and conformal blocks, Preprint (2002), 1–13, `math.QA/0210040`.

[MTV] Markov Y., Tarasov V., Varchenko A.: The determinant of a hypergeometric period matrix, Houston J. of Math. **24** (1998) No. 2, 197–220.

[MV] Markov Y., Varchenko A.: Hypergeometric solutions of trigonometric KZ equations satisfy dynamical difference equations, Adv. Math. **166** (2002), no. 1, 100–147.

[MuV] Mukhin E., Varchenko A.: The quantized Knizhnik-Zamolodchikov equation in tensor products of irreducible $sl(2)$-modules, Calogero-Moser-Sutherland models (Montreal, QC, 1997), CRM Series Math. Phys., Springer, New York, 2000.

[TL] Toledano-Laredo, V.: A Kohno-Drinfeld theorem for quantum Weyl groups, Duke Math. J. **112** (2002), no. 3 421–451.

[TV1] Tarasov, V., Varchenko, A.: Geometry of q-hypergeometric functions as a bridge between Yangians and quantum affine algebras, Invent. Math. **128** (1997), 501–588.

[TV2] Tarasov, V., Varchenko, A.: Difference equations compatible with trigonometric KZ differential equations, Int. Math. Res. Notices (2000), no. 15, 801-829.

[TV3] Tarasov, V., Varchenko, A.: Duality for Knizhnik-Zamolodchikov and dynamical equations, Acta Appl. Math. **73** (2002), no. 1-2, 141–154.

[TV4] Tarasov, V., Varchenko, A. : Identities for hypergeometric integrals of different dimensions, Preprint (2003), 1–9, `math.QA/0305224`.

[TV5] Tarasov, V., Varchenko, A.: Dynamical differential equations compatible with rational qKZ equations, Preprint (2004), 1–7, `math.QA/0403416`.

[TV6] Tarasov, V., Varchenko, A.: Hypergeometric solutions of rational qKZ equations satisfy dynamical differential equations, in preparation.
[TV7] Tarasov, V., Varchenko, A.: Selberg type integrals associated with $\mathfrak{sl}_3$, Lett. Math. Phys. **65** (2003), 173–185.